

- Aber, J.D. see Wessman, C.A., 6
- Adams, W.T. see Merkle, S.A., 181
- Adkins, C.W. see Nelson, R.M., Jr., 391
- Alaback, P.B., and Herman, F.R. Long-term response of under-story vegetation to stand density in *Picea-Tsuga* forests, 1522
- Alig, R.J. see Parks, P.J., 965
- Amman, G.D., McGregor, M.D., Schmitz, R.F., and Oakes, R.D. Susceptibility of lodgepole pine to infestation by mountain pine beetles following partial cutting of stands, 688
- Armentano, T.V. see Muir, P.S., 498
- Arnott, J.T., and Burdett, A.N. Early growth of planted western hemlock in relation to stock type and controlled-release fertilizer application, 710
- Arp, P.A., and Manasc, J. Red spruce stands downwind from a coal-burning power generator: tree-ring analysis, 251
- Arthaud, G.J., and Klemperer, W.D. Optimizing high and low thinning in loblolly pine with dynamic programming, 1118
- Aussenac, G., and Granier, A. Effects of thinning on water stress and growth in Douglas-fir, 100
- Ballard, L.A., and Long, J.N. Influence of stand density on log quality of lodgepole pine, 911
- Ballard, T.M. see Vihnanek, R.E., 718
- Bare, B.B., and Mendoza, G.A. A soft optimization approach to forest land management planning, 545
- Barrett, J. see Farmer, R.E., Jr., 1078
- Barry, R., Plamondon, A.P., and Stein, J. Hydrologic soil properties and application of a soil moisture model in a balsam fir forest, 427
- Bartel-Ortiz, L.M., and David, M.B. Sulfur constituents and transformations in upland and floodplain forest soils, 1106
- Bartram, C., and Miller, G. Estimation of seed orchard efficiencies by means of multistage variable probability sampling, 1397
- Basham, J.T. Decay and stain 10 years later in aspen suckers subjected to scarification at age 3, 1507
- Beatty, S.W., and Sholes, O.D.V. Leaf litter effect on plant species composition of deciduous forest treefall pits, 553
- Beckett, D.R. see Jamnick, M.S., 330
- Bedker, P.J., and Blanchette, R.A. Mortality of Scots pine following inoculation with the pinewood nematode, *Bursaphelenchus xylophilus*, 574
- Beeson, R. see Margolis, H.A., 962
- Beeson, R.C., Jr., and Proebsting, W.M. Photosynthate translocation during union development in *Picea* grafts, 986
- Bégin, J. see Margolis, H.A., 962
- Bell, J.C. see Moran, G.F., 506
- Bella, I.E., and Navratil, S. Western gall rust dynamics and impact in young lodgepole pine stands in west-central Alberta, 1437
- Bella, I.E. see Zakrzewski, W.T., 195
- Bella, I.E. see Tait, D.E., 1255
- Bellefleur, P. see Messier, C., 615
- Bellefleur, P. see Margolis, H.A., 962
- Bentzer, B.G., Foster, G.S., Hellberg, A.R., and Podzorski, A.C. Genotype \times environment interaction in Norway spruce involving three levels of genetic control: seed source, clone mixture, and clone, 1172
- Bergeron, J.-M., and Tardif, J. Winter browsing preferences of snowshoe hares for coniferous seedlings and its implication in large-scale reforestation programs, 280
- Bernier, B., and Brazeau, M. Foliar nutrient status in relation to sugar maple dieback and decline in the Quebec Appalachians, 754
- Bernier, B., and Brazeau, M. Nutrient deficiency symptoms associated with sugar maple dieback and decline in the Quebec Appalachians, 762
- Bernier, B., and Brazeau, M. Magnesium deficiency symptoms associated with sugar maple dieback in a Lower Laurentians site in southeastern Quebec, 1265
- Bernier, B., and Brazeau, M. An occurrence of boron deficiency in the deciduous forest of the Quebec Appalachians and the St. Lawrence Lowlands, 1652
- Berryman, A.A. see Lieutier, F., 1243
- Biging, G.S. Estimating the accuracy of volume equations using taper equations of stem profile, 1002
- Bingham, B.B., and Sawyer, J.O., Jr. Volume and mass of decaying logs in an upland old-growth redwood forest, 1649
- Binkley, C.S. see Cardellicchio, P.A., 981
- Binkley, D. see Hedman, C.W., 1090
- Black, T.A. see Livingston, N.J., 1234
- Blake, J.I., and Hoogenboom, G. A dynamic simulation of loblolly pine (*Pinus taeda* L.) seedling establishment based upon carbon and water balances, 833
- Blanchette, R.A. see Bedker, P.J., 574
- Blenis, P.V., and Pinnell, H.D. Effect of inoculum concentration of *Endocronartium harknessii* on the infection of lodgepole pine, 1123
- Blenis, P.V., Wiggins, K.L., Cunningham, J.E., and Pickard, M.A. Maltol protects against infection of lodgepole pine seedlings by western gall rust, 1658
- Bloomberg, W.J., and Brix, H. Effect of root injury and auxin treatments on adventitious root development in second-growth Douglas-fir, 957
- Blum, B.M. Variation in the phenology of bud flushing in white and red spruce, 315
- Blumen, A. see von Niessen, W., 805
- Bockheim, J.G. see Ruark, G.A., 435
- Boring, L.R. see White, D.L., 54
- Boyce, R.L. Wind direction and fir wave motion, 461
- Bozzuto, L.M., and Wilson, B.F. Branch angle in red maple trees, 643
- Brand, D.G., and Janas, P.S. Growth and acclimation of planted white pine and white spruce seedlings in response to environmental conditions, 320
- Brand, D.G., and Magnussen, S. Asymmetric, two-sided competition in even-aged monocultures of red pine, 901
- Brann, T.B. see Reams, G.A., 787
- Brazeau, M. see Bernier, B., 754
- Brazeau, M. see Bernier, B., 762
- Brazeau, M. see Bernier, B., 1265
- Brazeau, M. see Bernier, B., 1652
- Breuil, C., Seifert, K.A., Yamada, J., Rossignol, L., and Saddler, J.N. Quantitative estimation of fungal colonization of wood using an enzyme-linked immunosorbent assay, 374
- Brix, H. see Bloomberg, W.J., 957

- Brown, B.G., and Murphy, A.H.** On the economic value of weather forecasts in wildfire suppression mobilization decisions, 1641
- Brown, K.M., and Mugasha, A.G.** A two-stage method for horizontal point sampling in young forest stands, 1340
- Brown, K.R., Zobel, D.B., and Zasada, J.C.** Seed dispersal, seedling emergence, and early survival of *Larix laricina* (DuRoi) K. Koch in the Tanana Valley, Alaska, 306
- Brumelis, G., and Carleton, T.J.** The vegetation of postlogged black spruce lowlands in central Canada. I. Trees and tall shrubs, 1470
- Bunnell, F.L.** see Vales, D.J., 606
- Buongiorno, J., Chavas, J.-P., and Uusivuori, J.** Exchange rates, Canadian lumber imports, and United States prices: a time-series analysis, 1587
- Burdett, A.N.** see Amott, J.T., 710
- Burk, T.E.** see Sievänen, R., 1027
- Burkhart, H.E.** see Dyer, M.E., 825
- Calamassi, R., Falusi, M., and Mugnai, L.** Shoot morphology and growth pattern in seedlings of *Pinus brutia* provenances, 188
- Campbell, R.K.** see Merkle, S.A., 181
- Cardellicchio, P.A., and Binkley, C.S.** The effects of overrun improvements on stumpage price inflation, 981
- Carleton, T.J.** see Brumelis, G., 1470
- Carlson, C.E., McCaughey, W.W., and Theroux, L.J.** Relations among stand structure, dispersal of second-instar western spruce budworm, defoliation, and height growth of young conifers, 794
- Carlson, W.C., Harrington, C.A., Farnum, P., and Hallgren, S.W.** Effects of root severing treatments on loblolly pine, 1376
- Carmean, W.H.** see Schmidt, M.G., 297
- Carter, G.A., Smith, W.K., and Hadley, J.L.** Stomatal conductance in three conifer species at different elevations during summer in Wyoming, 242
- Castello, J.D.** see Ferris, M.A., 813
- Cates, R.G.** see Van Home, B., 90
- Cates, R.G.** see Paine, T.D., 1556
- Catling, P.M., and Spicer, K.W.** The separation of *Betula populifolia* and *Betula pendula* and their status in Ontario, 1017
- Ceulemans, R., Impens, I., and Steenackers, V.** Genetic variation in aspects of leaf growth of *Populus* clones, using the leaf plastochron index, 1069
- Chabot, M.** see Gagnon, R.R., 1655
- Chakravarty, P., and Chatarpaul, L.** The effects of Velpar L (hexazinone) on seedling growth and ectomycorrhizal symbiosis *Pinus resinosa*, 917
- Chatarpaul, L.** see Chakravarty, P., 917
- Chauret, G.** see Keith, C.T., 1325
- Chavas, J.-P.** see Buongiorno, J., 1587
- Cheliak, W.M., Wang, J., and Pitel, J.A.** Population structure and genic diversity in tamarack, *Larix laricina* (Du Roi) K. Koch, 1318
- Cheliak, W.M.** see Farmer, R.E., Jr., 1078
- Chong, N.** see Juzwik, J., 1493
- Cieszewski, C.J.** see Tait, D.E., 1255
- Clancy, K.M., Wagner, M.R., and Tinus, R.W.** Variation in host foliage nutrient concentrations in relation to western spruce budworm herbivory, 530
- Clatterbuck, W.K., and Hodges, J.D.** Development of cherrybark oak and sweet gum in mixed, even-aged bottomland stands in central Mississippi, U.S.A., 12
- Cleland, D.T.** see Host, G.E., 659
- Clements, S.E., Leuschner, W.A., Hoganson, H., and Wisdom, H.W.** A local roundwood supply and price assessment model, 1563
- Colombo, S.J.** see Koppelaar, R.S., 1103
- Conkey, L.E.** Decline in old-growth red spruce in western Maine: an analysis of wood density and climate, 1063
- Constantino, L.F.** Sawlog prices and quality differences in Canadian and United States Pacific coastal log markets, 540
- Constantino, L.F., and Haley, D.** Wood quality and the input and output choices of sawmilling producers for the British Columbia coast and the United States Pacific Northwest, west side, 202
- Cook, S.P., and Hain, F.P.** Wound response of loblolly and shortleaf pine attached or reattacked by *Dendrotonus frontalis* Zimmermann (Coleoptera: Scolytidae) or its fungal associate, *Ceratocystis minor* (Hedgecock) Hunt, 33
- Corns, I.G.W.** Compaction by forestry equipment and effects on coniferous seedling growth on four soils in the Alberta foothills, 75
- Courchesne, F., et Hendershot, W.H.** Cycle annuel des éléments nutritifs dans un bassin-versant forestier : contribution de la litière fraîche, 930
- Cregg, B.M., Dougherty, P.M., and Hennessey, T.C.** Growth and wood quality of young loblolly pine trees in relation to stand density and climatic factors, 851
- Critchfield, W.B.** see St. Clair, J.B., 640
- Cromack, K.** see Helgerson, O.T., 1082
- Cromack, K., Jr.** see Heath, B., 68
- Cronan, C.S.** see Rustad, L.E., 947
- Crow, T.R.** see Erdmann, G.G., 134
- Cunningham, J.E.** see Blenis, P.V., 1658
- Cwynar, L.C.** Late Quaternary vegetation history of Kettlehole Pond, southwestern Yukon, 1270
- Dancik, B.P.** see Govindaraju, D.R., 1347
- Dancik, B.P.** see Kenny, J.R., 1595
- David, M.B., Grigal, D.F., Ohmann, L.F., and Gertner, G.Z.** Sulfur, carbon, and nitrogen relationships in forest soils across the northern Great Lake States as affected by atmospheric deposition and vegetation, 1386
- David, M.B.** see Bartel-Ortiz, L.M., 1106
- DeBell, D.S.** see Franklin, J.F., 633
- Delisle, G.P., Woodard, P.M., Titus, S.J., and Johnson, A.F.** Sample size and variability of fuel weight estimates in natural stands of lodgepole pine, 649
- Dell, T.R.** see Van Deusen, P.C., 825
- Desprez-Loustau, M.L., et Dessureault, M.** Influence de stress contrôlés sur la sensibilité du bouleau jaune au chancre godronien causé par *Godronia cassandrae* Peck f.sp. *betulicola* Groves, 121
- Dessureault, M.** see Desprez-Loustau, M.L., 121
- DeYoe, D.R.** see Tung, C.-H., 1486
- Dictus, K.** see Godbold, D.L., 1167
- Diebolt, K.S., and Mudge, K.W.** Use of a video-imaging system for estimating leaf surface area of *Pinus sylvestris* seedlings, 377
- Dodd, R.S.** Cyclophytic effects on wood structure in *Pinus radiata* D. Don. II. Tracheid morphology, 413
- Dodd, R.S., and Walker, N.K.** Cyclophytic effects on wood structure in *Pinus radiata* D. Don. I. Densitometry and grain angle, 406
- Dougherty, P.M.** see Cregg, B.M., 851
- Drew, A.P.** Interference of black cherry by ground flora of the Allegheny uplands, 652
- Dumas, C.** see Villar, M., 1261
- Dumas, M.T.** Biological species of *Armillaria* in the mixedwood forest of northern Ontario, 872

- Dunstan, D.I. Prospects and progress in conifer biotechnology, 1497
- Dyer, M.E., and Burkhart, H.E. Reply: Compatible crown ratio and crown height models by P.C. Van Deusen and T.R. Dell, 825
- Eis, S. Erratum: Root systems of older immature hemlock, cedar, and Douglas-fir, 657
- Ek, A.R. see Sievänen, R., 1027
- Eldridge, K.G. see Moran, G.F., 506
- El-Kassaby, Y.A. see Thomson, A.J., 515
- Erdmann, G.G., Crow, T.R., and Rauscher, H.M. Foliar nutrient variation and sampling intensity for *Acer rubrum* trees, 134
- Fahey, T.J., Yavitt, J.B., and Joyce, G. Precipitation and throughfall chemistry in *Pinus contorta* ssp. *latifolia* ecosystems, southeastern Wyoming, 337
- Fall, M. see Iles, K., 774
- Falusi, M. see Calamassi, R., 188
- Farmer, R.E., Jr., Garlick, K., and Watson, S.R. Heritability and C effects in a 3-year-old balsam poplar clonal test, 1059
- Farmer, R.E., Jr., Cheliak, W.M., Perry, D.J., Knowles, P., Barrett, J., and Pitel, J.A. Isozyme variation in balsam poplar along a latitudinal transect in northwestern Ontario, 1078
- Farnum, P. see Carlson, W.C., 1376
- Farrar, R.M., Jr. see Murphy, P.A., 827
- Federer, C.A. see Hornbeck, J.W., 1337
- Fedosejevs, G. see Leckie, D.G., 1008
- Feller, M.C. see Trowbridge, R., 128
- Fenn, R.C. see Stettler, R.F., 745
- Feret, P.P. see Johnsen, K.H., 610
- Ferm, A. see Kauppi, A., 1603
- Ferris, M.A., and Castello, J.D. Detection of tomato ringspot virus in white ash and adjacent vegetation in central New York, 813
- Fites, J.A., and Teskey, R.O. CO₂ and water vapor exchange of *Pinus taeda* in relation to stomatal behavior: test of an optimization hypothesis, 150
- Fitzgerald, J.W. see Watwood, M.E., 820
- Flinn, B.S. see Webb, D.T., 1570
- Florence, L.Z. see Kenny, J.R., 1595
- Fortin, J.A. see Gagnon, J., 922
- Foster, G.S. see Bentzer, B.G., 1172
- Fowler, D.P., Park, Y.S., and Gordon, A.G. Genetic variation of red spruce in the Maritimes, 703
- Fowler, D.P. see Park, Y.S., 106
- Franklin, J.F., and DeBell, D.S. Thirty-six years of tree population change in an old-growth *Pseudotsuga-Tsuga* forest, 633
- Fried, J.S., Tappeiner, J.C., II, and Hibbs, D.E. Bigleaf maple seedling establishment and early growth in Douglas-fir forests, 1226
- Fullerton, J.M. see Martell, D.L., 444
- Gaget, M. see Villar, M., 1261
- Gagnon, J., Langlois, C.G., and Fortin, J.A. Growth and ectomycorrhiza formation of containerized black spruce seedlings as affected by nitrogen fertilization, inoculum type, and symbiont, 922
- Gagnon, R.R., Chabot, M., et Pineau, M. Effets de l'élagage sur la résistance électrique cambiale de la tige de jeunes Sapins baumiers, 1655
- Gagnon, R.R. see Margolis, H.A., 723
- Garlick, K. see Farmer, R.E., Jr., 1059
- Geiger, J.P. see Nandris, D., 1248
- Georgis, W. see Webb, D.T., 1570
- Gerhold, H.D. see Stanton, B.J., 1531
- Geron, C.D., and Ruark, G.A. Comparison of constant and variable allometric ratios for predicting foliar biomass of various tree genera, 1298
- Gertner, G.Z. see David, M.B., 1386
- Gillard, D. see Tóth, J., 353
- Gjerstad, D.H. see Kuhns, M.R., 285
- Glerum, C. see Kim, Y.T., 1286
- Glover, G.R. see Williams, H.M., 1635
- Godbold, D.L., Dictus, K., and Hüttermann, A. Influence of aluminium and nitrate on root growth and mineral nutrition of Norway spruce (*Picea abies*) seedlings, 1167
- Goheen, D.J. see Hansen, E.M., 942
- Golden, M.S. see Tuttle, C.L., 628
- Golden, M.S. see Tuttle, C.L., 867
- Gonzalez, J.S., and Richards, J. Early selection for wood density in young coastal Douglas-fir trees, 1182
- Gordon, A.G. see Fowler, D.P., 703
- Govindaraju, D.R., Wagner, D.B., Smith, G.P., and Dancik, B.P. Chloroplast DNA variation within individual trees of a *Pinus banksiana* — *Pinus contorta* sympatric region, 1347
- Grace, J.C. see Santantonio, D., 657
- Granier, A. see Aussenac, G., 100
- Gregoire, T.G., and Walters, D.K. Composite vector estimators derived by weighting inversely proportional to variance, 282
- Grier, C.C. Foliage loss due to snow, wind, and winter drying damage: its effects on leaf biomass of some western conifer forests, 1097
- Grigal, D.F. see David, M.B., 1386
- Grönroos, R., and von Arnold, S. Initiation of roots on hypocotyl cuttings of *Pinus sylvestris*, with emphasis on direct rooting, root elongation, and auxin uptake, 1457
- Groot, A. Methods for estimating seedbed receptivity and for predicting seedling stocking and density in broadcast seeding, 1541
- Hadley, J.L. see Carter, G.A., 242
- Hain, F.P. see Cook, S.P., 33
- Haines, B.L. see White, D.L., 54
- Haley, D. see Constantino, L.F., 202
- Hall, D.J. see Raffa, K.F., 1661
- Hallam, P.M., and Tibbitts, W.N. Determination of frost hardness of *Eucalyptus* using the electrical conductivity of diffusate in conjunction with a freezing chamber, 595
- Hallgren, S.W., and Helms, J.A. Control of height growth components in seedlings of California red and white fir by seed source and water stress, 521
- Hallgren, S.W. see Carlson, W.C., 1376
- Halteman, W.A. see Reams, G.A., 787
- Hamm, P.B. see Hansen, E.M., 1053
- Hamrick, J.L. see Smith, C.C., 453
- Hänell, B. Postdrainage forest productivity of peatlands in Sweden, 1443
- Hanley, T.A. see Van Horne, B., 90
- Hänninen, H., and Pelkonen, P. Does the temperature response of rest break of woody plants change during the development? 269
- Hänninen, H., and Pelkonen, P. Erratum: Does the temperature response of rest break of woody plants change during the development? 1214
- Hansen, E.A., McLaughlin, R.A., and Pope, P.E. Biomass and nitrogen dynamics of hybrid poplar on two different soils: implications for fertilization strategy, 223
- Hansen, E.M., and Hamm, P.B. Canker diseases of Douglas-fir seedlings in Oregon and Washington bareroot nurseries, 1053

- Hansen, E.M., and Goheen, D.J. Rate of increase of black-stain root disease in Douglas-fir plantations in Oregon and Washington, 942
- Harrington, C.A. see Carlson, W.C., 1376
- Harrington, T.C. see Rizzo, D.M., 991
- Harrington, T.C. see Worral, J.J., 1463
- Heath, B., Sollins, P., Perry, D.A., and Cromack, K., Jr. Asymbiotic nitrogen fixation in litter from Pacific Northwest forests, 68
- Hedman, C.W., and Binkley, D. Canopy profiles of some Piedmont hardwood forests, 1090
- Heilman, P.E. see Stettler, R.F., 745
- Helgeson, O.T., Cromack, K., Stafford, S., Miller, R.E., and Slagle, R. Equations for estimating aboveground components of young Douglas-fir and red alder in a coastal Oregon plantation, 1082
- Hellberg, A.R. see Bentzer, B.G., 1172
- Helms, J.A. see Hallgren, S.W., 521
- Hendershot, W.H. see Courchesne, F., 930
- Hendrickson, O.Q. Biomass and nutrients in regenerating woody vegetation following whole-tree and conventional harvest in a northern mixed forest, 1427
- Hennessey, T.C., Lorenzi, E.M., and McNew, R.W. Stomatal conductance and growth of five *Alnus glutinosa* clones in response to controlled water stress, 421
- Hennessey, T.C. see Cregg, B.M., 851
- Herman, F.R. see Alaback, P.B., 1522
- Hibbs, D.E. see Fried, J.S., 1226
- Hinckley, T.M. see Pezeshki, S.R., 1159
- Hiratsuka, Y. see Mallett, K.I., 292
- Hix, D.M. Multifactor classification and analysis of upland hardwood forest ecosystems of the Kickapoo River watershed, southwestern Wisconsin, 1405
- Ho, R.H. Gibberellin A₄₇ enhances seed-cone production in field-grown black spruce, 139
- Hodges, J.D. see Clatterbuck, W.K., 12
- Hoganson, H. see Clements, S.E., 1563
- Honhart, C. see Juzwik, J., 1493
- Hoogenboom, G. see Blake, J.I., 833
- Hornbeck, J.W., Smith, R.B., and Federer, C.A. Growth trends in 10 species of trees in New England, 1950–1980, 1337
- Horner, J.D. see Van Horne, B., 90
- Host, G.E., Pregitzer, K.S., Ramm, C.W., Lusch, D.P., and Cleland, D.T. Variation in overstory biomass among glacial landforms and ecological land units in northwestern Lower Michigan, 659
- Houston, D.R., and Valentine, H.T. Beech bark disease: the temporal pattern of cankering in aftermath forests of Maine, 38
- Howard, A.F. Harvesting profitability as affected by classification of costs and assignment of the components of scheduled time, 1369
- Hüttermann, A. see Godbold, D.L., 1167
- Iles, K., and Fall, M. Can an angle gauge really evaluate "borderline trees" accurately in variable plot sampling? 774
- Iles, K., and Wilson, W.H. Changing angle gauges in variable plot sampling: Is there a bias under ordinary conditions? 768
- Iles, K. see Smith, N.J., 467
- Impens, I. see Ceulemans, R., 1069
- Innes, J.L. Forest health surveys: problems in assessing observer objectivity, 560
- Jacobson, J.S. see Van Ryn, D.M., 1049
- Jahraus, K.V. see Tait, D.E., 1479
- Jamnick, M.S., and Beckett, D.R. A logit analysis of private woodlot owner's harvesting decisions in New Brunswick, 330
- Janas, P.S. see Brand, D.G., 320
- Jin, L., van der Kamp, B.J., Wilson, J., and Swan, E.P. Biodegradation of thujaplicins in living western red cedar, 782
- Johnsen, K.H., Feret, P.P., and Seiler, J.R. Root growth potential and shoot activity of northern and southern provenances of 1–0 eastern white pine seedlings grown in a Virginia nursery, 610
- Johnson, A.F. see Delisle, G.P., 649
- Jokela, J.J. see Uddin, M.R., 937
- Joslin, J.D., and Wolfe, M.H. Responses of red spruce seedlings to changes in soil aluminum in six amended forest soil horizons, 1614
- Joyce, D.G. Adaptive variation in cold hardness of eastern larch, *Larix laricina*, in northern Ontario, 85
- Joyce, G. see Fahey, T.J., 337
- Jozsa, L.A. see Robertson, E.O., 888
- Juzwik, J., Honhart, C., and Chong, N. Cylindrocladium root rot in Ontario bare-root nurseries: estimate of spruce seedling losses, 1493
- Kauppi, A., Kiviniitty, M., and Ferm, A. Growth habits and crown architecture of *Betula pubescens* Ehrh. of seed and sprout origin, 1603
- Keifer, M.B. see Kimball, K.D., 385
- Keith, C.T., and Chauret, G. Basic wood properties of European larch from fast-growth plantations in eastern Canada, 1325
- Kenney, W.A. see Mosseler, A., 1535
- Kenny, J.R., Dancik, B.P., Florence, L.Z., and Nargang, F.E. Nucleotide sequence of the carboxy-terminal portion of a lodgepole pine actin gene, 1595
- Kim, Y.T., and Glerum, C. Free amino acid concentrations in red pine needles during three successive autumns, 1286
- Kimball, K.D., and Keifer, M. Climatic comparisons with tree-ring data from montane forests: are the climatic data appropriate? 385
- Kiss, G., and Yeh, F.C. Heritability estimates for height for young interior spruce in British Columbia, 158
- Kittredge, D.B., Jr. The influence of species composition on the growth of individual red oaks in mixed stands in southern New England, 1550
- Kiviniitty, M. see Kauppi, A., 1603
- Klemperer, W.D. see Arthaud, G.J., 1118
- Knowles, P. see Farmer, R.E., Jr., 1078
- Koppelaar, R.S., and Colombo, S.J. Heat tolerance of actively growing, bud-initiated, and dormant black spruce seedlings, 1103
- Korelus, V.J. see Thomson, A.J., 1343
- Kozak, A. A variable-exponent taper equation, 1363
- Kramer, C.L. see Smith, C.C., 453
- Kubler, H. Silvicultural control of mechanical stresses in trees, 1215
- Kuhns, M.R., and Gjerstad, D.H. Photosynthate allocation in loblolly pine (*Pinus taeda*) seedlings as affected by moisture stress, 285
- Kyker-Snowman, T.D., and Wilson, B.F. Total wood, sapwood, and heartwood in branch bases of three conifers, 1332
- Langlois, C.G. see Gagnon, J., 922
- La Roi, G.H., Strong, W.L., and Pluth, D.J. Understory plant community classifications as predictors of forest site quality for lodgepole pine and white spruce in west-central Alberta, 875
- Lassoie, J.P. see Van Ryn, D.M., 1049
- Lavigne, M.B. Stem growth and respiration of young balsam fir trees in thinned and unthinned stands, 483
- Lavigne, M.B. Growth and net assimilation rates in thinned and unthinned stands of balsam fir, 1205
- Leckie, D.G., Teillet, P.M., Fedosejevs, G., and Ostaff, D.P.

- Reflectance characteristics of cumulative defoliation of balsam fir, 1008
- Leininger, T.D., and Winner, W.E. Throughfall chemistry beneath *Quercus rubra*: atmospheric, foliar, and soil chemistry considerations, 478
- Leuschner, W.A. see Clements, S.E., 1563
- Li, H.G. see Schreuder, H.T., 1280
- Lieutier, F., and Berryman, A.A. Preliminary histological investigations of the defense reactions of three pines to *Ceratocystis clavigera* and two chemical elicitors, 1243
- Lineberger, R.D. see Struve, D.K., 265
- Livingston, N.J., and Black, T.A. The growth and water use of three species of conifer seedlings planted on a high-elevation south-facing clearcut, 1234
- Loehle, C. Tree life history strategies: the role of defenses, 209
- Long, J.N., and Smith, F.W. Leaf area – sapwood area relations of lodgepole pine as influenced by stand density and site index, 247
- Long, J.N. see Ballard, L.A., 911
- Lorenzi, E.M. see Hennessey, T.C., 421
- Loustau, D. see Ruel, J.-C., 1196
- Lusch, D.P. see Host, G.E., 659
- MacDougall, R.G., MacLean, D.A., and Thompson, R.G. The use of electrical capacitance to determine growth and vigor of spruce and fir trees and stands in New Brunswick, 587
- MacLean, D.A. see MacDougall, R.G., 587
- MacMillan, P.C. Decomposition of coarse woody debris in an old-growth Indiana forest, 1353
- Magnussen, S., and Yeatman, C.W. Height growth and survival of within- and between-provenance crosses in jack pine, 1145
- Magnussen, S. see Brand, D.G., 901
- Malcolm, D.C. see McKay, H.M., 1416
- Mallett, K.I., and Hiratsuka, Y. Inoculation studies of lodgepole pine with Alberta isolates of the *Armillaria mellea* complex, 292
- Manasc, J. see Arp, P.A., 251
- Margolis, H.A., Gagnon, R.R., Pothier, D., and Pineau, M. The adjustment of growth, sapwood area, heartwood area, and sapwood saturated permeability of balsam fir after different intensities of pruning, 723
- Margolis, H.A., Bégin, J., Beeson, R., and Bellefleur, P. The influence of metal halide and high-pressure sodium lamps during photoperiod extension on the allocation of carbon between lignin and cellulose in black spruce seedlings, 962
- Marshall, P.L. A procedure for constructing timber management strategies under uncertainty, 398
- Martell, D.L., and Fullerton, J.M. Decision analysis for jack pine management, 444
- McCarthy, B.C. A method of access into the crowns of subcanopy and canopy trees, 646
- McCaughy, J.H. see Saxton, W.L., 143
- McCaughy, W.W. see Carlson, C.E., 794
- McGregor, M.D. see Amman, G.D., 688
- McKay, H.M., and Malcolm, D.C. A comparison of the fine root component of a pure and a mixed coniferous stand, 1416
- McKendrick, J.D. see Van Horne, B., 90
- McLaughlin, R.A. see Hansen, E.A., 223
- McLean, J.A., and Tuytel, J. Marking forest insects: evaluation of two systems for the systemic introduction of rubidium into Douglas-fir trees, 19
- McLean, J.A. see Shore, T.L., 1113
- McLeod, K.W. see Topa, M.A., 276
- McLeod, S.D., and Running, S.W. Comparing site quality indices and productivity in ponderosa pine stands of western Montana, 346
- McMinn, J.W., and Nutter, W.L. Season and intensity of whole-tree harvesting influence regeneration in the oak–pine type, 669
- McNay, R.S., Peterson, L.D., and Nyberg, J.B. The influence of forest stand characteristics on snow interception in the coastal forests of British Columbia, 566
- McNew, R.W. see Hennessey, T.C., 421
- Meil, J.K., and Nautiyal, J.C. An intraregional economic analysis of production structure and factor demand in major Canadian softwood lumber producing regions, 1036
- Meldahl, R.S. see Tuttle, C.L., 628
- Meldahl, R.S. see Tuttle, C.L., 867
- Melillo, J.M. see Wessman, C.A., 6
- Mendoza, G.A. see Bare, B.B., 545
- Merkle, S.A., Adams, W.T., and Campbell, R.K. Multivariate analysis of allozyme variation patterns in coastal Douglas-fir from southwest Oregon, 181
- Messier, C., and Bellefleur, P. Light quantity and quality on the forest floor of pioneer and climax stages in a birch–beech–sugar maple stand, 615
- Meyer, M.M., Jr. see Uddin, M.R., 937
- Micko, M.M. see Yang, R.C., 954
- Miller, G. see Bartram, C., 1397
- Miller, R.E. see Helgeson, O.T., 1082
- Misson, J.-P. Multiplication du *Thuja plicata* par culture *in vitro* de tissus juvéniles et âgés, 473
- Moran, G.F., Bell, J.C., and Eldridge, K.G. The genetic structure and the conservation of the five natural populations of *Pinus radiata*, 506
- Mosseler, A., Zsuffa, L., Stoehr, M.U., and Kenney, W.A. Variation in biomass production, moisture content, and specific gravity in some North American willows (*Salix* L.), 1535
- Mudge, K.W. see Diebolt, K.S., 377
- Mugasha, A.G. see Brown, K.M., 1340
- Mugnai, L. see Calamassi, R., 188
- Muir, P.S., and Armentano, T.V. Evaluating oxidant injury to foliage of *Pinus ponderosa*: a comparison of methods, 498
- Murphy, A.H. see Brown, B.G., 1641
- Murphy, P.A., and Farrar, R.M., Jr. Basal-area projection equations for thinned natural even-aged forest stands, 827
- Nandris, D., Nicole, M., and Geiger, J.P. Root-rot diseases of the rubber tree in the Ivory Coast. 1. Severity, dynamics, and characterization of epidemics, 1248
- Nargang, F.E. see Kenny, J.R., 1595
- Nautiyal, J.C. see Meil, J.K., 1036
- Navratil, S. see Bella, I.E., 1437
- Nelson, E.E. see Thies, W.G., 801
- Nelson, R.M., Jr., and Adkins, C.W. A dimensionless correlation for the spread of wind-driven fires, 391
- Newnham, R.M. A modification of the Ek-Payandeh nonlinear regression model for site index curves, 115
- Nicole, M. see Nandris, D., 1248
- Nutter, W.L. see McMinn, J.W., 669
- Nyberg, J.B. see McNay, R.S., 566
- Oakes, R.D. see Amman, G.D., 688
- O'Hara, K.L. Stand structure and growing space efficiency following thinning in an even-aged Douglas-fir stand, 859
- Ohmann, L.F. see David, M.B., 1386
- O'Reilly, C., and Owens, J.N. Reproductive growth and development in seven provenances of lodgepole pine, 43
- Ostaf, D.P. see Leckie, D.G., 1008
- Ouellet, D., and Zarnovican, R. Cultural treatment of young yellow birch (*Betula alleghaniensis* Britton) stands. 1. Tree classification and stand structure, 1581

- Ouellet, D. see Ung, C.-H., 739
- Owens, J.N., and Simpson, S.J. Bud and shoot development in *Picea engelmannii* in response to cone induction treatments, 231
- Owens, J.N. see O'Reilly, C., 43
- Paine, T.D., Stephen, F.M., and Cates, R.G. Phenology of an induced response in loblolly pine following inoculation of fungi associated with the southern pine beetle, 1556
- Pallardy, S.G. see Parker, W.C., 1
- Pallardy, S.G. see Parker, W.C., 1211
- Park, Y.S., and Fowler, D.P. Geographic variation of black spruce tested in the Maritimes, 106
- Park, Y.S. see Fowler, D.P., 703
- Parker, W.C., and Pallardy, S.G. Leaf and root osmotic adjustment in drought-stressed *Quercus alba*, *Q. macrocarpa*, and *Q. stellata* seedlings, 1
- Parker, W.C., and Pallardy, S.G. Pressure-volume analysis of leaves of *Robinia pseudoacacia* L. with the sap expression and free transpiration methods, 1211
- Parker, W.H., van Niejenhuis, A., and Van Damme, L. Baseline selection of black spruce by large-scale aerial photography, 380
- Parks, P.J., and Alig, R.J. Land base models for forest resource supply analysis: a critical review, 965
- Payandeh, B. Discussion: An assessment of the structural method of deriving a black spruce site equation by V.G. Smith, 1351
- Pelkonen, P. see Hänninen, H., 269
- Pelkonen, P. see Hänninen, H., 1214
- Perry, D.A. see Heath, B., 68
- Perry, D.J. see Farmer, R.E., Jr., 1078
- Peterson, D.L. see Wessman, C.A., 6
- Peterson, L.D. see McNay, R.S., 566
- Peterson, M.J., Sutherland, J.R., and Tuller, S.E. Greenhouse environment and epidemiology of grey mould of container-grown Douglas-fir seedlings, 974
- Pezeshki, S.R., and Hinckley, T.M. Water relations characteristics of *Alnus rubra* and *Populus trichocarpa*: responses to field drought, 1159
- Phipps, R.L., and Whiton, J.C. Decline in long-term growth trends of white oak, 24
- Pickard, M.A. see Blenis, P.V., 1658
- Pineau, M. see Margolis, H.A., 723
- Pineau, M. see Ruel, J.-C., 1196
- Pineau, M. see Gagnon, R.R., 1655
- Pinnell, H.D. see Blenis, P.V., 1123
- Pitel, J.A. see Farmer, R.E., Jr., 1078
- Pitel, J.A. see Cheliak, W.M., 1318
- Plamondon, A.P. see Barry, R., 427
- Pluth, D.J. see La Roi, G.H., 875
- Podzorski, A.C. see Bentzer, B.G., 1172
- Pope, P.E. see Hansen, E.A., 223
- Pope, P.E. see Simmons, G.L., 728
- Pope, P.E. see Simmons, G.L., 1392
- Pothier, D. see Margolis, H.A., 723
- Pregitzer, K.S. see Host, G.E., 659
- Price, A.G. see Watters, R.J., 1490
- Proebsting, W.M. see Beeson, R.C., Jr., 986
- Raffa, K.F., and Hall, D.J. *Thrips calcaratus* Uzel (Thysanoptera: Thripidae), a new pest of basswood trees in the Great Lakes region, 1661
- Raffa, K.F., and Smalley, E.B. Response of red and jack pines to inoculation with microbial associates of the pine engraver, *Ips pini* (Coleoptera: Scolytidae), 581
- Raffa, K.F., and Smalley, E.B. Host resistance to invasion by lower stem and root infesting insects of pine: response to controlled inoculations with the fungal associate *Leptographium terebrantis*, 675
- Raffa, K.F., and Smalley, E.B. Seasonal and long-term responses of host trees to microbial associates of the pine engraver, *Ips pini*, 1624
- Ralph, D.R. see Thomson, A.J., 1343
- Ramm, C.W. see Host, G.E., 659
- Rapacher, H.M. see Erdmann, G.G., 134
- Reams, G.A., Brann, T.B., and Halteman, W.A. A nonparametric survival model for balsam fir during a spruce budworm outbreak, 787
- Reinhardt, E.D. see Ryan, K.C., 1291
- Rhodus, W.T. see Struve, D.K., 131
- Richards, J. see Gonzalez, J.S., 1182
- Rizzo, D.M., and Harrington, T.C. Root movement and root damage of red spruce and balsam fir on subalpine sites in the White Mountains, New Hampshire, 991
- Robertson, E.O., and Jozsa, L.A. Climatic reconstruction from tree rings at Banff, 888
- Ross, S.D. Effects of temperature, drought, and gibberellin A₄/7, and timing of treatment, on flowering in potted *Picea engelmannii* and *Picea glauca* grafts, 163
- Ross, S.D. Pre- and post-pollination polyhouse environment effects on pollen and seed development in potted *Picea engelmannii* grafts, 623
- Rossignol, L. see Breuil, C., 374
- Ruark, G.A., and Bockheim, J.G. Biomass, net primary production, and nutrient distribution for an age sequence of *Populus tremuloides* ecosystems, 435
- Ruark, G.A. see Geron, C.D., 1298
- Ruel, J.-C., Loustau, D., et Pineau, M. Relations entre la microtopographie, les caractéristiques de la couverture morte et la répartition des essences dans une érablière à Bouleau jaune, 1196
- Running, S.W. see McLeod, S.D., 346
- Rustad, L.E., and Cronan, C.S. Element loss and retention during litter decay in a red spruce stand in Maine, 947
- Ryan, K.C., and Reinhardt, E.D. Predicting postfire mortality of seven western conifers, 1291
- Ryan, M.G. see Schreuder, H.T., 1280
- Saddler, J.N. see Breuil, C., 374
- Santantonio, D., and Grace, J.C. Erratum: Estimating fine-root production and turnover from biomass and decomposition data: a compartment-flow model, 657
- Sawyer, J.O., Jr. see Bingham, B.B., 1649
- Saxton, W.L., and McCaughey, J.H. Measurement considerations and trends in biomass heat storage of a mixed forest, 143
- Schmidt, M.G., and Carmean, W.H. Jack pine site quality in relation to soil and topography in north central Ontario, 297
- Schmitz, R.F. see Amman, G.D., 688
- Schreuder, H.T., Li, H.G., Ryan, M.G., and Scott, C.T. Adjusting estimates in two-way tables by incorporating outside information, 1280
- Scott, C.T. see Schreuder, H.T., 1280
- Seifert, K.A. see Breuil, C., 374
- Seiler, J.R. see Johnsen, K.H., 610
- Shay, J.M. see Zwiazek, J.J., 1305
- Shay, J.M. see Zwiazek, J.J., 1311
- Sholes, O.D.V. see Beatty, S.W., 553
- Shore, T.L., and McLean, J.A. The use of mark-recapture to evaluate a pheromone-based mass trapping program for ambrosia beetles in a sawmill, 1113
- Sievänen, R., Burk, T.E., and Ek, A.R. Construction of a stand

- growth model utilizing photosynthesis and respiration relationships in individual trees, 1027
- Simmons, G.L., and Pope, P.E.** Development of a root growth model for yellow-poplar and sweetgum seedlings grown in compacted soil, 728
- Simmons, G.L., and Pope, P.E.** Influence of soil water potential and mycorrhizal colonization on root growth of yellow-poplar and sweet gum seedlings grown in compacted soil, 1392
- Simpson, S.J.** see Owens, J.N., 231
- Slagle, R.** see Helgerson, O.T., 1082
- Smalley, E.B.** see Raffa, K.F., 581
- Smalley, E.B.** see Raffa, K.F., 675
- Smalley, E.B.** see Raffa, K.F., 1624
- Smit, B.A.** Selection of flood-resistant and susceptible seedlings of *Populus trichocarpa* Torr. & Gray, 271
- Smith, C.C., Hamrick, J.L., and Kramer, C.L.** The effects of stand density on frequency of filled seeds and fecundity in lodgepole pine (*Pinus contorta* Dougl.), 453
- Smith, F.W.** see Long, J.N., 247
- Smith, G.P.** see Govindaraju, D.R., 1347
- Smith, N.J., and Iles, K.** A graphical depiction of multivariate similarity among sample plots, 467
- Smith, R.B.** see Hornbeck, J.W., 1337
- Smith, V.G.** Reply: An assessment of the structural method of deriving a black spruce site equation by B. Payandeh, 1351
- Smith, W.K.** see Carter, G.A., 242
- Sollins, P.** see Heath, B., 68
- South, D.B.** see Tuttle, C.L., 867
- South, D.B.** see Williams, H.M., 1635
- Spicer, K.W.** see Catling, P.M., 1017
- Stafford, S.** see Helgerson, O.T., 1082
- Stanton, B.J., and Gerhold, H.D.** Family and family \times nitrogen interaction effects on juvenile growth of *Prunus serotina*, 1531
- Stanton, B.J.** see Stettler, R.F., 745
- St. Clair, J.B., and Critchfield, W.B.** Hybridization of a Rocky Mountain fir (*Abies concolor*) and a Mexican fir (*Abies religiosa*), 640
- Steenackers, V.** see Ceulemans, R., 1069
- Stein, J.** see Barry, R., 427
- Stephen, F.M.** see Paine, T.D., 1556
- Stettler, R.F., Fenn, R.C., Heilman, P.E., and Stanton, B.J.** *Populus trichocarpa* \times *Populus deltoides* hybrids for short rotation culture: variation patterns and 4-year field performance, 745
- Stoehr, M.U.** see Mosseler, A., 1535
- Stohlgren, T.J.** Litter dynamics in two Sierran mixed conifer forests. I. Litterfall and decomposition rates, 1127
- Stohlgren, T.J.** Litter dynamics in two Sierran mixed conifer forests. II. Nutrient release in decomposing leaf litter, 1136
- Strong, W.L.** see La Roi, G.H., 875
- Struve, D.K., and Lineberger, R.D.** Restoration of high adventitious root regeneration potential in mature *Betula papyrifera* Marsh. softwood stem cuttings, 265
- Struve, D.K., and Rhodus, W.T.** Phenyl indole-3-thiolobutyrate increases growth of transplanted 1-0 red oak, 131
- Sutherland, J.R.** see Peterson, M.J., 974
- Swan, E.P.** see Jin, L., 782
- Swank, W.T.** see Watwood, M.E., 820
- Tait, D.E.** The dynamics of stand development: a general stand model applied to Douglas-fir, 696
- Tait, D.E., and Jahraus, K.V.** The contrast in stand dynamics as revealed by comparing parameter estimates for a general stand growth model, 1479
- Tait, D.E., Cieszewski, C.J., and Bella, I.E.** The stand dynamics of lodgepole pine, 1255
- Tappeiner, J.C., II.** see Fried, J.S., 1226
- Tardif, J.** see Bergeron, J.-M., 280
- Teillet, P.M.** see Leckie, D.G., 1008
- Teskey, R.O.** see Fites, J.A., 150
- Theroux, L.J.** see Carlson, C.E., 794
- Thies, W.G., and Nelson, E.E.** Bulldozing stumps and nitrogen fertilization affect growth of Douglas-fir seedlings, 801
- Thompson, R.G.** see MacDougall, R.G., 587
- Thomson, A.J., and El-Kassaby, Y.A.** Trend surface analysis of a Douglas-fir provenance-progeny transfer test, 515
- Thomson, A.J., Tudor, K.D., Korelus, V.J., and Ralph, D.R.** Detecting the response of Douglas-fir to nitrogen fertilization by regression of periodic annual basal area increment against basal area, 1343
- Tibbits, W.N.** see Hallam, P.M., 595
- Tinus, R.W.** see Clancy, K.M., 530
- Titus, S.J.** see Delisle, G.P., 649
- Todd, D.** The effects of host genotype, growth rate, and needle age on the distribution of a mutualistic, endophytic fungus in Douglas-fir plantations, 601
- Topa, M.A., and McLeod, K.W.** Promotion of aerenchyma formation in *Pinus serotina* seedlings by ethylene, 276
- Tóth, J., and Gillard, D.** Experimental design and evaluation of a peatland drainage system for forestry by optimization of synthetic hydrographs, 353
- Trowbridge, R., and Feller, M.C.** Relationships between the moisture content of the fine woody fuels in lodgepole pine slash and the Fine Fuel Moisture Code of the Canadian Forest Fire Weather Index System, 128
- Tudor, K.D.** see Thomson, A.J., 1343
- Tuller, S.E.** see Peterson, M.J., 974
- Tung, C.-H., and DeYoe, D.R.** Growth resumption and performance of nonchilled *Abies* seedlings after their first winter, 1486
- Tuttle, C.L., Golden, M.S., and Meldahl, R.S.** Soil compaction effects on *Pinus taeda* establishment from seed and early growth, 628
- Tuttle, C.L., South, D.B., Golden, M.S., and Meldahl, R.S.** Initial *Pinus taeda* seedling height relationships with early survival and growth, 867
- Tuytel, J.** see McLean, J.A., 19
- Uddin, M.R., Meyer, M.M., Jr., and Jokela, J.J.** Plantlet production from anthers of Eastern cottonwood (*Populus deltoides*), 937
- Ung, C.-H., et Végiard, S.** Problèmes d'inférence statistique reliés à la transformation logarithmique en régression, 733
- Ung, C.-H., Végiard, S., et Ouellet, D.** Variance asymptotique des estimateurs en régression non linéaire, 739
- Uusivuori, J.** see Buongiorno, J., 1587
- Valentine, H.T.** see Houston, D.R., 38
- Vales, D.J., and Bunnell, F.L.** Comparison of methods for estimating forest overstory cover. I. Observer effects, 606
- Van Damme, L.** see Parker, W.H., 380
- van den Driessche, R.** Nursery growth of conifer seedlings using fertilizers of different solubilities and application time, and their forest growth, 172
- van der Kamp, B.J.** Susceptibility of lodgepole pine provenances to geographically separate western gall rust spore sources, 1203
- van der Kamp, B.J.** see Jin, L., 782
- Van Deusen, P.C.** Simultaneous estimation with a squared error loss function, 1093
- Van Deusen, P.C., and Dell, T.R.** Discussion: Compatible crown

- ratio and crown height models by M.E. Dyer and H.E. Burkhart, 825
- Van Horne, B., Hanley, T.A., Cates, R.G., McKendrick, J.D., and Horner, J.D. Influence of seral stage and season on leaf chemistry of southeastern Alaska deer forage, 90
- van Niejenhuis, A. see Parker, W.H., 380
- Van Ryn, D.M., Lassoie, J.P., and Jacobson, J.S. Effects of acid mist on *in vivo* pollen tube growth in red maple, 1049
- Van Wagner, C.E. Effect of slope on fires spreading downhill, 818
- Végiard, S. see Ung, C.-H., 733
- Végiard, S. see Ung, C.-H., 739
- Vihnanek, R.E., and Ballard, T.M. Slashburning effects on stocking, growth, and nutrition of young Douglas-fir plantations in salal-dominated ecosystems of eastern Vancouver Island, 718
- Villar, M., Gaget, M., and Dumas, C. Micro-isoelectric focusing of proteins from single stigmas of *Populus*, 1261
- Volney, W.J.A. Analysis of historic jack pine budworm outbreaks in the Prairie provinces of Canada, 1152
- von Arnold, S. see Grönroos, R., 1457
- von Niessen, W., and Blumen, A. Dynamic simulation of forest fires, 805
- Wagner, D.B. see Govindaraju, D.R., 1347
- Wagner, M.R. see Clancy, K.M., 530
- Walker, N.K. see Dodd, R.S., 406
- Wall, R.E. Deterioration of severely defoliated balsam fir in relation to stand age, spacing, and foliar protection, 490
- Walters, D.K. see Gregoire, T.G., 282
- Wang, E.I.C. see Yang, R.C., 954
- Wang, J. see Cheliak, W.M., 1318
- Watson, S.R. see Farmer, R.E., Jr., 1059
- Watters, R.J., and Price, A.G. The influence of stemflow from standing dead trees on the fluxes of some ions in a mixed deciduous forest, 1490
- Watwood, M.E., Fitzgerald, J.W., and Swank, W.T. Effects of moisture content on sulfate generation and retention in hardwood forest upper soil horizons, 820
- Webb, D.T., Flinn, B.S., and Georgis, W. Micropropagation of eastern white pine (*Pinus strobus* L.), 1570
- Webb, S.L. Windstorm damage and microsite colonization in two Minnesota forests, 1186
- Wessman, C.A., Aber, J.D., Peterson, D.L., and Melillo, J.M. Foliar analysis using near infrared reflectance spectroscopy, 6
- White, D.L., Haines, B.L., and Boring, L.R. Litter decomposition in southern Appalachian black locust and pine-hardwood stands: litter quality and nitrogen dynamics, 54
- Whiton, J.C. see Phipps, R.L., 24
- Wiggins, K.L. see Blenis, P.V., 1658
- Williams, C.G. Accelerated short-term genetic testing for loblolly pine families, 1085
- Williams, H.M., South, D.B., and Glover, G.R. Effect of bud status and seedling biomass on root growth potential of loblolly pine, 1635
- Wilson, A.A.G. Width of firebreak that is necessary to stop grass fires: some field experiments, 682
- Wilson, B.F. see Bozzuto, L.M., 643
- Wilson, B.F. see Kyker-Snowman, T.D., 1332
- Wilson, J. see Jin, L., 782
- Wilson, W.H. see Iles, K., 768
- Winner, W.E. see Leininger, T.D., 478
- Wisdom, H.W. see Clements, S.E., 1563
- Wolfe, M.H. see Joslin, J.D., 1614
- Woodard, P.M. see Delisle, G.P., 649
- Worrall, J.J., and Harrington, T.C. Etiology of canopy gaps in spruce-fir forests at Crawford Notch, New Hampshire, 1463
- Yamada, J. see Breuil, C., 374
- Yang, R.C., Wang, E.I.C., and Micko, M.M. Effects of fertilization on wood density and tracheid length of 70-year-old lodgepole pine in west-central Alberta, 954
- Yavitt, J.B. see Fahey, T.J., 337
- Yeatman, C.W. see Magnussen, S., 1145
- Yeh, F.C. see Kiss, G., 158
- Zakrzewski, W.T., and Bella, I.E. Two new height models for volume estimation of lodgepole pine stands, 195
- Zarnovican, R. see Ouellet, D., 1581
- Zasada, J.C. Embryo growth in Alaskan white spruce seeds, 64
- Zasada, J.C. see Brown, K.R., 306
- Zobel, D.B. see Brown, K.R., 306
- Zsuffa, L. see Mosseler, A., 1535
- Zwiazek, J.J., and Shay, J.M. Sodium fluoride induced metabolic changes in jack pine seedlings. I. Effect on gas exchange, water content, and carbohydrates, 1305
- Zwiazek, J.J., and Shay, J.M. Sodium fluoride induced metabolic changes in jack pine seedlings. II. Effect on growth, acid phosphatase, cytokinins, and pools of soluble proteins, amino acids, and organic acids, 1311

